

[View as a Web page](#)



Introducing a New Treatment Option for Previously Treated Recurrent or Metastatic Cervical Cancer

Dear Ms. Chinchilla,

We are sending this communication on behalf of Seagen, to announce the approval of Tivdak™ (tisotumab vedotin-tftv) for injection 40 mg, for intravenous use, by the U.S. Food and Drug Administration (FDA) on September 20, 2021. Tivdak is a tissue factor-directed antibody and microtubule inhibitor conjugate indicated for the treatment of adult patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy.

This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

Please see Important Safety Information below and the full [Prescribing Information, including BOXED WARNING](#).¹

Product Name	Strength	NDC-11 Code ²	Wholesale Acquisition Cost (WAC)
Tivdak (tisotumab vedotin-tftv)	40 mg single-dose vial	51144-0003-01*	\$5,885

*The NDC has been "zero-filled" to ensure creation of an 11-digit code that meets Health Insurance Portability and Accountability Act (HIPAA) standards. The zero-fill location is indicated in bold.

IMPORTANT SAFETY INFORMATION

BOXED WARNING: OCULAR TOXICITY

TIVDAK caused changes in the corneal epithelium and conjunctiva resulting in changes in vision, including severe vision loss, and corneal ulceration. Conduct an ophthalmic exam at baseline, prior to each dose, and as clinically indicated. Adhere to premedication and required eye care before, during, and after infusion. Withhold TIVDAK until improvement and resume, reduce the dose, or permanently discontinue, based on severity.

Warnings and Precautions

Ocular Adverse Reactions occurred in 60% of patients with cervical cancer treated with TIVDAK across clinical trials. The most common were conjunctival adverse reactions (40%), dry eye (29%), corneal adverse reactions (21%), and blepharitis (8%). Grade 3 ocular adverse reactions occurred in 3.8% of patients, including severe ulcerative keratitis in 3.2% of patients. One patient experienced ulcerative keratitis with perforation requiring corneal transplantation. Cases of symblepharon were reported in patients with other tumor types treated with TIVDAK at the recommended dose.

In innovaTV 204, 4% of patients experienced visual acuity changes to 20/50 or worse including 1% of patients who experienced a visual acuity change to 20/200. Of the patients who experienced decreased visual acuity to 20/50 or worse, 75% resolved, including the patient who experienced decreased visual acuity to 20/200.

Refer patients to an eye care provider for an ophthalmic exam including visual acuity and slit lamp exam at baseline, prior to each dose, and as clinically indicated. Adhere to premedication and required eye care to reduce the risk of ocular adverse reactions. Promptly refer patients to an eye care provider for any new or worsening ocular signs and symptoms. Withhold dose, reduce the dose, or permanently discontinue TIVDAK based on the severity of the adverse reaction.

Peripheral Neuropathy (PN) occurred in 42% of cervical cancer patients treated with TIVDAK across clinical trials; 8% of patients experienced Grade 3 PN. PN adverse reactions included peripheral neuropathy (20%), peripheral sensory neuropathy (11%), peripheral sensorimotor neuropathy (5%), motor neuropathy (3%), muscular weakness (3%), and demyelinating peripheral polyneuropathy (1%). One patient with another tumor type treated with TIVDAK at the recommended dose developed Guillain-Barre syndrome. Monitor patients for signs and symptoms of neuropathy. For new or worsening PN, withhold, dose reduce, or permanently discontinue TIVDAK based on the severity of PN.

Hemorrhage occurred in 62% of cervical cancer patients treated with TIVDAK across clinical trials. The most common all grade hemorrhage adverse

reactions were epistaxis (44%), hematuria (10%), and vaginal hemorrhage (10%). Grade 3 hemorrhage occurred in 5% of patients.

Monitor patients for signs and symptoms of hemorrhage. For patients experiencing pulmonary or CNS hemorrhage, permanently discontinue TIVDAK. For Grade ≥ 2 hemorrhage in any other location, withhold until bleeding has resolved, blood hemoglobin is stable, there is no bleeding diathesis that could increase the risk of continuing therapy, and there is no anatomical or pathologic condition that can increase the risk of hemorrhage recurrence. After resolution, either resume treatment or permanently discontinue TIVDAK.

Pneumonitis: Severe, life-threatening, or fatal pneumonitis can occur in patients treated with antibody-drug conjugates containing vedotin, including TIVDAK. Among patients with cervical cancer treated with TIVDAK across clinical trials, 2 patients (1.3%) experienced pneumonitis, including 1 patient who had a fatal outcome.

Monitor patients for pulmonary symptoms of pneumonitis. Infectious, neoplastic, and other causes for symptoms should be excluded through appropriate investigations.

Withhold TIVDAK for patients who develop persistent or recurrent Grade 2 pneumonitis and consider dose reduction. Permanently discontinue TIVDAK in all patients with Grade 3 or 4 pneumonitis.

Embryo-Fetal Toxicity: TIVDAK can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TIVDAK and for 2 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with TIVDAK and for 4 months after the last dose.

Adverse Reactions

Serious adverse reactions occurred in 43% of patients; the most common ($\geq 3\%$) were ileus (6%), hemorrhage (5%), pneumonia (4%), PN, sepsis, constipation, and pyrexia (each 3%). Fatal adverse reactions occurred in 4% of patients who received TIVDAK, including septic shock, pneumonitis, sudden death, and multisystem organ failure (each 1%).

Adverse reactions leading to permanent discontinuation occurred in 13% of patients receiving TIVDAK; the most common ($\geq 3\%$) were PN (5%) and corneal adverse reactions (4%). Adverse reactions leading to dose interruption occurred in 47% of patients; the most common ($\geq 3\%$) were PN (8%), conjunctival adverse reactions (4%), and hemorrhage (4%). Adverse reactions leading to dose reduction occurred in 23% of patients; the most common ($\geq 3\%$) were conjunctival adverse reactions (9%) and corneal adverse reactions (8%).

The most common ($\geq 25\%$) adverse reactions, including laboratory abnormalities, were hemoglobin decreased (52%), fatigue (50%), lymphocytes decreased (42%), nausea (41%), PN (39%), alopecia (39%), epistaxis (39%), conjunctival adverse reactions (37%), hemorrhage (32%), leukocytes decreased (30%), creatinine increased (29%), dry eye (29%), prothrombin international normalized ratio increased (26%), activated partial thromboplastin time prolonged (26%), diarrhea (25%), and rash (25%).

Drug interactions

Strong CYP3A4 Inhibitors: Concomitant use with strong CYP3A4 inhibitors may increase unconjugated monomethyl auristatin E (MMAE) exposure, which may increase the risk of TIVDAK adverse reactions. Closely monitor patients for TIVDAK adverse reactions.

Use in Specific Populations

Moderate or Severe Hepatic Impairment: MMAE exposure and adverse reactions are increased. Avoid use.

Lactation: Advise lactating women not to breastfeed during TIVDAK treatment and for at least 3 weeks after the last dose.

For additional information about Tivdak, please see the full [Prescribing Information including BOXED WARNING](#).

Please visit <http://www.tivdakhcp.com> to learn more about Tivdak. For questions regarding Tivdak, please contact Seagen Medical Information at 1-855-4SEAGEN (1-855-473-2436) or medinfo@seagen.com.

Sincerely,

Aaron Franczek
Consultant
DK Pierce and Associates, Inc.
10910 Creek Way
Zionsville, IN 46077
(317) 873-0303
aaron.franczek@dkpierce.net

References

1. TIVDAK™ (tisotumab vedotin-tftv) [Prescribing Information]. Bothell, WA: Seagen Inc. September 2021.
2. US Food and Drug Administration. National Drug Code Background Information. <https://www.fda.gov/drugs/development-approval-process-drugs/national-drug-code-database-background-information>. Accessed June 22, 2021.

Tivdak and its logo are trademarks of Seagen Inc. Seagen and its logo are US registered trademarks of Seagen Inc. Bothell, WA 98021.

© 2021 Seagen Inc. and Genmab. All rights reserved. US-TVP-21-188-MT

[Unsubscribe](#) | [Privacy Policy](#)

This email was sent to: carmencg@nebmed.org

This email was sent by: **DK Pierce & Associates, Inc.**
10910 Creek Way
Zionsville, IN 46077

(317) 873-0303

